

## 1 Dihedral groups

Suppose that we have numbered the vertices of a regular  $n$ -gon by  $\{1, 2, \dots, n\}$ . Notice that there are exactly  $n$  choices to replace the first vertex. If we replace the first vertex by  $k$ , then the second vertex must be replaced either by vertex  $k + 1$  or by vertex  $k - 1$  (to be a rigid motion); hence, there are at most  $2n$  possible rigid motions of the regular  $n$ -gon. The group of symmetries of the regular  $n$ -polygon is denoted by  $\mathbb{D}_n$ .

Let us denote, for  $k = 0, 1, \dots, n - 1$ , by  $r_k$  the counter-clock rotation with angle

$$\theta_k = \frac{360^\circ k}{n}.$$

Also, for  $k = 1, 2, \dots, n$ , we denote by  $s_k$ , the reflexion around the axis of symmetry through vertex  $k$ . If  $k$  is even there are only  $n/2$  such different reflexions. On the other hand if  $k$  is odd, there will be  $n$  such reflexions. The reflexions  $s_k$  satisfy  $s_k^2 = 1$  and the rotation  $r_k^n = 1$ . We denote  $s_1 = s$  and  $r_1 = r$ .

**Lemma 1.** *The elements  $r, s$  satisfy the relation  $sr^j = r^{-j}s$ .*

*Proof.* Let  $s$  be the reflection around the axis of symmetry through vertex 1. Take any vertex  $k$  and consider the action of both maps on  $k \bmod n$ :

$$sr^j(k) \equiv s(k + j) \equiv 2 - k - j \equiv r^{-j}(2 - k) \equiv r^{-j}s(k).$$

And this would take care of the relation we wanted to prove. □

**Theorem 2.** *The group  $\mathbb{D}_n$ , with  $n \geq 3$ , consists of all products of the two elements  $r$  and  $s$ , satisfying the relations*

$$r^n = 1, \quad s^2 = 1 \quad \text{and} \quad srs = r^{-1}.$$

*Proof.* Any rigid motion  $t$  of the  $n$ -gon replacing the first vertex by the vertex  $k$ , must replace the second vertex by an adjacent vertex to  $k$ . If the second vertex goes to  $k + 1$ , then  $t = r^k$ . If the second vertex is replaced by  $k - 1$ , then  $t = r^k s$ . Hence,  $r$  and  $s$  generate  $\mathbb{D}_n$  and

$$\mathbb{D}_n = \{1, r, r^2, \dots, r^{n-1}, s, rs, \dots, r^{n-1}s\},$$

by lemma 1. □

**Remark 3.** The rotation  $r_k$  admits a matrix representation:

$$r_k = \begin{pmatrix} \cos\left(\frac{360^\circ k}{n}\right) & -\sin\left(\frac{360^\circ k}{n}\right) \\ \sin\left(\frac{360^\circ k}{n}\right) & \cos\left(\frac{360^\circ k}{n}\right) \end{pmatrix}$$

We can check that the matrices that we obtain in this case are elements of  $\text{Sl}_2(\mathbb{R})$ . In general we say that we have a representation of a group  $G$  when we have a good map (group homomorphism)

$$\rho: G \longrightarrow \text{Gl}(V),$$

for a vector space  $V$  over  $\mathbb{C}$ . When the dimension of  $V$  is 1, this particular case of representation is called a character.

### Practice Questions:

1. Find a 2-dimensional matrix representation for the reflexions of  $\mathbb{D}_n$ .