

**Math 32, Homework 3 on sections 5.3, 6.1, 6.2**

---

Do these 10 questions and *check that your answers match the solutions on page 2*. They will not be collected, but similar questions could appear on the next quiz.

- (1) Sketch the region enclosed by the following curves and lines

$$y = x^2, \quad y = 0, \quad x = 1, \quad x = 2.$$

Then use the method of cylindrical shells to find the volume of the solid obtained by rotating this region about the  $y$ -axis.

- (2) Sketch the region enclosed by the following curve and line

$$y = 4x - x^2, \quad y = x.$$

Then use the method of cylindrical shells to set up the integral to find the volume of the solid obtained by rotating this region about the  $y$ -axis. (No need to compute the integral.)

- (3) Is the function  $f(x) = 2 + \cos(x)$  one-to-one? Is it one-to-one on the domain  $0 \leq x \leq \pi$ ?  
(4) Let  $g(x) = 3x + 1$ . Find  $g^{-1}(x)$  and graph both  $g$  and its inverse together.

- (5) Let

$$f(x) = \frac{x + 1}{2x - 3}.$$

Find the inverse of  $f$ . Then give the domain and range of  $f$  and the domain and range of  $f^{-1}$ .

- (6) (a) Use the formula you found in Q5 to compute  $(f^{-1})'(1)$ . (b) Use the formula

$$(f^{-1})'(x) = \frac{1}{f'(f^{-1}(x))}$$

to compute  $(f^{-1})'(1)$ .

- (7) Graph the exponential function  $h(x) = (0.4)^x$ . Give the domain and range of  $h$  and state whether it is increasing or decreasing. Also find  $\lim_{x \rightarrow \infty} h(x)$  and  $\lim_{x \rightarrow -\infty} h(x)$ .

- (8) Find:

$$(a) \quad \lim_{x \rightarrow \infty} \frac{2e^x - 1}{e^x + 1}, \quad (b) \quad \lim_{x \rightarrow \infty} e^{-x} \sin(x).$$

- (9) Differentiate:

$$(a) \quad e^{4 \tan(x)}, \quad (b) \quad e^{x/(x-1)}.$$

(10) Differentiate:

$$(a) \sqrt{1 + e^{x^2}}, \quad (b) 3x^2 e^{\cos(x)}.$$

---

You can also try questions from sections 5.3, 6.1, 6.2 in the book listed on the syllabus.

---

**Answers to questions (1)-(10):**

(1) The volume is  $15\pi/2$ .

(2) The integral giving the volume is

$$2\pi \int_0^3 (3x^2 - x^3) dx.$$

(3) With the horizontal line test,  $f(x)$  is not one-to-one. But it is one-to-one on the domain  $0 \leq x \leq \pi$ .

(4)  $g^{-1}(x) = (x - 1)/3$ . The graphs of  $g$  and  $g^{-1}$  are symmetric about the line  $y = x$ .

(5) We have

$$f^{-1}(x) = \frac{3x + 1}{2x - 1}$$

and

$$\begin{aligned} \text{domain of } f &= \mathbb{R}/\{3/2\}, & \text{range of } f &= \mathbb{R}/\{1/2\} \\ \text{domain of } f^{-1} &= \mathbb{R}/\{1/2\}, & \text{range of } f^{-1} &= \mathbb{R}/\{3/2\}. \end{aligned}$$

(6) With both methods  $(f^{-1})'(1) = -5$ .

(7) We have

$$\text{domain of } h = \mathbb{R}, \quad \text{range of } h = (0, \infty).$$

This function is decreasing and  $\lim_{x \rightarrow \infty} h(x) = 0$ ,  $\lim_{x \rightarrow -\infty} h(x) = \infty$ .

(8) We have

$$(a) \lim_{x \rightarrow \infty} \frac{2e^x - 1}{e^x + 1} = 2, \quad (b) \lim_{x \rightarrow \infty} e^{-x} \sin(x) = 0.$$

(9) We have

$$(a) \frac{d}{dx} e^{4 \tan(x)} = 4 \sec^2(x) e^{4 \tan(x)}, \quad (b) \frac{d}{dx} e^{x/(x-1)} = \frac{-1}{(x-1)^2} e^{x/(x-1)}.$$

(10) We have

$$(a) \frac{d}{dx} \sqrt{1 + e^{x^2}} = \frac{x e^{x^2}}{\sqrt{1 + e^{x^2}}}, \quad (b) \frac{d}{dx} 3x^2 e^{\cos(x)} = 3x(2 - x \sin(x)) e^{\cos(x)}.$$