

# 1 Characters

## 1.1 Group characters for finite abelian groups

Characters will play a central role in the classification of finite abelian groups. Consider the set  $C = \{z \in \mathbb{C}^* \mid |z| = 1\}$  in the complex plane.

**Proposition 1.**  $C$  is a subgroup of  $\mathbb{C}^* = \{z \in \mathbb{C} \mid z \neq 0\}$  and  $\xi^{-1} = \bar{\xi}$  (complex conjugate).

*Proof.* The identity element 1 of  $\mathbb{C}^*$  is in  $C$ . Also for  $z_1, z_2 \in \mathbb{C}$  we have  $|z_1 z_2| = |z_1| |z_2|$ , and the set  $C$  is closed under multiplications. For the inverse we have:

$$z \in C \Rightarrow z^{-1} = \frac{1}{z} = \frac{\bar{z}}{|z|^2} = \bar{z} \in C.$$

And we have checked all conditions for  $C$  being a subgroup. □

**Proposition 2.** Every finite subgroup of  $C$  is cyclic.

*Proof.* Let  $D$  be a subgroup of  $C$ . If  $D = 1$  then  $D = \langle 1 \rangle$  and we are done. Otherwise, suppose that  $D \neq 1$  and choose the element  $\delta \in D$  such that the positive angle  $\theta$  to the  $x$ -axis is minimal. If the element  $z \in D$  is strictly  $\delta^k < z < \delta^{k+1}$  for some value  $k$ , then the element  $z\delta^{-k} \in D$  would have a smaller angle. The conclusion is that, for any  $z \in D$  there exists a power  $k$  such that  $z = \delta^k$ . □

**Example 3.** The group  $C_n$  of complex  $n$ -roots of unity is a finite subgroup of  $C$ .

**Remark 4.** Infinite subgroups of  $C$  do not need to be cyclic. Take for example an irrational number  $\theta$  and the group  $K$  generated by  $-1$  and  $e^{2\pi i\theta}$ . If this group were to be cyclic generated by an element  $\delta = e^{2\pi i\phi}$ , we will have integers  $k, l$  such that

$$e^{2\pi i\theta} = \delta^k = e^{2k\pi i\phi} \text{ and } -1 = \delta^l = e^{2l\pi i\phi}.$$

This will give  $\theta = k\phi + m$  and  $\frac{1}{2} = l\phi + n$  for integers  $m, n$ . Since,  $\theta$  is irrational and  $1/2$  is a rational number we arrive to a contradiction. In the group just described we cannot define an element with minimum positive angle.

**Definition 5.** Let  $G$  be a finite abelian group. A character of  $G$  is a group homomorphism  $\chi: G \rightarrow \mathbb{C}^*$ . The set of all characters of  $G$  is denoted by  $\widehat{G}$ .

**Proposition 6.** The set of characters has a natural structure of abelian group.

*Proof.* Define  $\chi * \chi'$  by the map multiplication  $(\chi * \chi')(x) = \chi(x)\chi'(x)$ . It is also a group homomorphism  $G \rightarrow C$ . Also put  $\chi^{-1} = \bar{\chi}$  and take the identity element as the constant function 1.  $\square$

**Remark 7.** The group of characters of a finite abelian group is finite. Let  $x \in G$  and  $n$  be the order of the group  $G$ . We have  $1 = \chi(1) = \chi(x^n) = (\chi(x))^n$ . Hence  $\chi(x)$  is an  $n$ -th root of unity in  $C$ , there are at most  $n$  choices of  $\chi(x)$  for each  $x \in G$  and the number of characters is finite.

**Proposition 8.** *If  $G$  is cyclic,  $\widehat{G} \cong G$ .*

*Proof.* Let  $\chi$  be a character on  $G$  and  $G = \langle g \rangle$  of order  $n$ . Since  $\chi(g)^n = 1$ , we know that  $\chi(g)$  is a complex  $n$ -root of unity, that is  $\chi(g) \in C_n$ . Let us define a map  $\rho: C_n \rightarrow \widehat{G}$ ,  $\rho(\xi) = \chi_\xi$ , where  $\chi_\xi(g^k) = \xi^k$ .

First of all, the map is well defined since  $g^k = g^{k'} \Rightarrow n|k - k' \Rightarrow \xi^k = \xi^{k'}$ . We need to check that:  $\chi_\xi$  is a character:  $\chi_\xi(g^k g^l) = \chi_\xi(g^{k+l}) = \xi^{k+l} = \xi^k \xi^l = \chi_\xi(g^k) \chi_\xi(g^l)$ .

The map is a group homomorphism:  $\chi_{\xi\nu}(g^k) = \xi^k \nu^k = \chi_\xi(g^k) \chi_\nu(g^k)$ .

The map is surjective: since each character  $\chi = \chi_\xi$  for some  $\xi \in C_n$ .

The map is injective: since  $\chi_\xi = 1 \Rightarrow \xi^k = 1$  for all  $k \Rightarrow \xi = 1$ .  $\square$

**Proposition 9.** *Let  $G$  be a finite abelian group and  $H \subset G$  a subgroup. Every character  $\chi_0$  on  $H$  can be extended to a character on  $G$ .*

*Proof.* We proceed by induction on the order of the quotient group  $|G/H|$ . If  $|G/H| = 1$ , then  $G = H$ , the character  $\chi_0$  is already a character of  $G$ . If the order of the quotient  $|G/H| > 1$ , choose a class  $gH$  of  $G/H$  such that  $gH \neq H$  and denote by  $M/H = \langle gH \rangle$  the cyclic group of  $G/H$  generated by  $gH$ . If  $M \neq G$ , both  $|M/H|$  and  $|G/M|$  are both of order less than  $|G/H|$  and hence we can extend  $\chi_0$ , first to a character of  $M$  and then to a character of the whole  $G$ . We may assume then that  $M = G$  and  $G/H = M/H = \langle gH \rangle$  is cyclic. Let  $n = |G/H|$ . We have then that  $g^n H = H$  and hence  $g^n \in H$ . Choose a complex number  $\xi$  such that  $\chi_0(g^n) = \xi^n$  and define the map:

$$\chi(g^r h) = \xi^r \chi_0(h),$$

for all  $n > r \geq 0$  and  $h \in H$ . The map is well defined because the cosets

$$H_r = \{g^r h \mid 0 \leq r < n, h \in H\}$$

form a partition of  $G$ . We can prove that the map defined this way is a character on the whole  $G$  and it extends  $\chi_0$ .  $\square$

**Corollary 10.** *Let  $G$  be a finite abelian group and  $h \in G$  such that  $h \neq 1$ . Then, there exist a character  $\chi \in \widehat{G}$  such that  $\chi(h) \neq 1$ .*

**Theorem 11.** *(Orthogonality relations) Let  $G$  be a finite abelian group:*

1. For  $\chi$  and  $\chi'$  characters we have  $\sum_{g \in G} \overline{\chi(g)} \chi'(g) = \begin{cases} |G| & \text{if } \chi = \chi' \\ 0 & \text{if } \chi \neq \chi' \end{cases}$

2. For  $g$  and  $g'$  in  $G$  have  $\sum_{\chi \in \hat{G}} \overline{\chi(g)} \chi(g') = \begin{cases} |\hat{G}| & \text{if } g = g' \\ 0 & \text{if } g \neq g' \end{cases}$

*Proof.* Consider the character  $\xi = \overline{\chi} \chi'$ . If  $\xi \equiv 1$ , then  $\xi(g) = 1$  for all  $g$  and  $\sum_{g \in G} \xi(g) = |G|$ . On the other hand if  $\xi$  is not identically 1, there exist  $g_1 \in G$  such that  $\xi(g_1) \neq 1$  and we will have:

$$\xi(g_1) \sum_{g \in G} \xi(g) = \sum_{g \in G} \xi(g_1 g) = \sum_{g \in G} \xi(g).$$

Since  $\xi(g_1) \neq 1$ , it must be the case that  $\sum_{g \in G} \xi(g) = 0$ .

The other direction is similar: put now  $h = g^{-1} g'$  and compute  $\sum_{\chi \in \hat{G}} \chi(h)$ . If  $h = 1$ , we get  $\sum_{\chi \in \hat{G}} \chi(h) = |\hat{G}|$ . On the other hand if  $h \neq 1$ , there is a character  $\chi_1(h) \neq 1$  and

$$\chi_1(h) \sum_{\chi \in \hat{G}} \chi(h) = \sum_{\chi \in \hat{G}} \chi_1(h) \chi(h) = \sum_{\chi \in \hat{G}} (\chi_1 \chi)(h) = \sum_{\chi \in \hat{G}} \chi(h)$$

and since  $\chi_1(h) \neq 1$  it must be the case that  $\sum_{\chi \in \hat{G}} \chi(h) = 0$ . □

**Theorem 12.** For each finite abelian group the orders  $|\hat{G}| = |G|$ .

*Proof.* We do a summation in two different ways:

$$|\hat{G}| + 0 \cdots + 0 = \sum_{g \in G} \sum_{\chi \in \hat{G}} \chi(g) = \sum_{\chi \in \hat{G}} \sum_{g \in G} \chi(g) = |G| + 0 + \cdots + 0$$

to get equality of the orders. □

## 1.2 Character tables

The characters of a finite group form a character table which encodes much useful information about the group  $G$ . Each row is labelled by a character and the entries in the row are the values of that character on elements of the group. The columns are labelled by elements of  $G$  (in a more general setting this would be representative of conjugacy classes). It is customary to label the first row by the trivial character  $\chi_0$ , which is the trivial action of  $G$  given by  $\rho(g) = 1$  for all  $g \in G$ .

	$C_0 = 1$	$C_1 = g_1$	$C_2 = g_2$	...
$\chi_0 = 1$	1	1	1	...
$\chi_1$	1	$\chi_1(g_1)$	$\chi_1(g_2)$	...
$\chi_2$	1	$\chi_2(g_1)$	$\chi_2(g_2)$	...
...	...	...	...	...

For example, suppose that  $\omega$  is a primitive third root of unity. Then the character table of the cyclic group  $C_3$  can be represented by:

	1	$g$	$g^2$
$\chi_0 = 1$	1	1	1
$\chi_1$	1	$\omega$	$\omega^2$
$\chi_2$	1	$\omega^2$	$\omega$

**Practice Questions:**

1. Describe the table character for the cyclic group  $C_5$ .